

Supplementary Table 5: Comparison of different responses in clinical scenarios according to practicing years

Clinical scenarios	Total cohort, n/number of responses (%)	<5 years, n/number of responses (%)	5-10 years, n/number of responses (%)	>10 years, n/number of responses (%)
TBI				
In 24 h without a repeated CT head	4/65 (6.15)	1/27 (3.7)	2/13 (15.38)	1/25 (4)
In 24 h if a repeated CT head is stable	26/65 (40)	10/27 (37.04)	6/13 (46.15)	10/25 (40)
In 48 h without a repeated CT head	3/65 (4.62)	1/27 (3.7)	0/13 (0)	2/25 (8)
In 48 h if a repeated CT head is stable	17/65 (26.15)	7/27 (25.93)	4/13 (30.77)	6/25 (24)
In 72 h without a repeated CT head	1/65 (1.54)	0/27 (0)	0/13 (0)	1/25 (4)
In 72 h if a repeated CT head is stable	12/65 (18.46)	8/27 (29.63)	1/13 (7.69)	3/25 (12)
TBI with ICP monitor				
In 24 h without a repeated CT head	7/65 (10.77)	2/27 (7.41)	1/13 (7.69)	4/25 (16)
In 24 h if a repeated CT head is stable	26/65 (40)	12/27 (44.44)	9/13 (69.23)	5/25 (20)
In 48 h without a repeated CT head	5/65 (7.70)	2/27 (7.41)	0/13 (0)	3/25 (12)
In 48 h if a repeated CT head is stable	10/65 (15.38)	3/27 (11.11)	2/13 (15.38)	5/25 (20)
In 72 h without a repeated CT head	7/65 (10.77)	2/27 (7.41)	1/13 (7.69)	4/25 (16)
In 72 h if a repeated CT head is stable	3/65 (4.62)	1/27 (3.7)	0/13 (0)	2/25 (8)
Wait until the ICP is within normal	1/65 (1.54)	1/27 (3.7)	0/13 (0)	0/25 (0)
Wait until the ventriculostomy is removed	6/65 (9.23)	4/27 (14.81)	0/13 (0)	2/25 (8)
SCI				
In 24 h without a repeated CT spine	21/65 (32.31)	10/27 (37.04)	5/13 (38.46)	6/25 (24)
In 24 h if a repeated CT spine is stable	15/65 (23.07)	7/27 (25.93)	3/13 (23.08)	5/25 (20)
In 48 h without a repeated CT spine	3/65 (4.62)	1/27 (3.70)	0/13 (0)	2/25 (8)
In 48 h if a repeated CT spine is stable	13/65 (20)	6/27 (22.22)	1/13 (7.69)	6/25 (24)
In 72 h without a repeated CT spine	2/65 (3.08)	0/27 (0)	2/13 (15.38)	0/25 (0)
In 72 h if a repeated CT spine is stable	4/65 (6.15)	0/27 (0)	1/13 (7.69)	3/25 (12)
NOR solid organ injuries: GSW through the liver				
In 24 h without a repeated CT abdomen	13/65 (20)	7/27 (25.93)	3/13 (23.08)	3/25 (12)
In 24 h if a repeated CT abdomen is stable	10/65 (15.38)	4/27 (14.81)	4/13 (30.77)	2/25 (8)
In 48 h without a repeated CT abdomen	15/65 (23.07)	8/27 (29.63)	2/13 (15.38)	5/25 (20)
In 48 h if a repeated CT abdomen is stable	7/65 (10.77)	3/27 (11.11)	2/13 (15.38)	2/25 (8)
In 72 h without a repeated CT abdomen	4/65 (6.15)	0/27 (0)	1/13 (7.69)	3/25 (12)
In 72 h if a repeated CT abdomen is stable	8/65 (12.31)	3/27 (11.11)	0/13 (0)	5/25 (20)
NOR solid organ injuries: Grade IV splenic injury				
In 24 h without a repeated CT abdomen	9/64 (14.06)	4/27 (14.81)	3/13 (23.08)	2/24 (8.33)
In 24 h if a repeated CT abdomen is stable	10/64 (15.63)	4/27 (14.81)	5/13 (38.46)	1/24 (4.17)
In 48 h without a repeated CT abdomen	11/64 (17.18)	5/27 (18.52)	2/13 (15.38)	4/24 (16.67)
In 48 h if a repeated CT abdomen is stable	8/64 (12.5)	3/27 (11.11)	1/13 (7.69)	4/24 (16.67)
In 72 h without a repeated CT abdomen	5/64 (7.81)	2/27 (7.41)	1/13 (7.69)	2/24 (8.33)
In 72 h if a repeated CT abdomen is stable	13/64 (20.31)	4/27 (14.81)	0/13 (0)	9/24 (37.50)

CT: Computed tomography, TBI: Traumatic brain injury, SCI: Spinal cord injury, NOR: Nonoperative, GSW: Gunshot wound, ICP: Intracranial pressure